arXiv:1704.04004v2 [gr-qc] 17 Apr 2017

Canonical gravity and scalar fields

Avadhut Purohit

Gogate Jogalekar College, Ratnagiri
University of Mumbai, India
avdhoot.purohit@gmail.com

April 18, 2017

Abstract

A combined variable theory of gravity and scalar field is developed under ADM formulation by redefining new canonical conjugate variables (Φ, Π). It is a dynamical theory of the space-matter with implicit time dependence. Quantizing this new field theory results in quantization of geometry and scalar field combined.

1 Introduction

•Motivation:

General Relativity and Quantum Field Theory are two most successful theories of 19^{th} century. General Relativity is a classical theory of dynamical space-time. Quantum theory tells us that every dynamical field has quantum properties. But General Relativity being a non-renormalizable theory is very difficult to quantize. Whereas, Quantum Field Theory is built on a fixed background and relies on Poincaré invariance. It is a quantum theory of matter fields. But General Relativity tells us that space-time is also dynamical. Therefore, both theories are still incomplete.

Anorwitt, Deser and Misner [1] formulated a theory which brings gravity closer to Lorentz covariant field theories by disentangling dynamical degrees of freedom from the gauge variables. Loop Quantum Gravity is one of the most promising amongst canonical theories of gravity. It is a dynamical theory of connections ([2] and [3] can be referred for further details). A granularity of space is a result of quantization program. But the theory is far from being a complete theory for various reasons such as the existence of semi-classical limit is not proven so far ([4] section VI, A summarizes some more open problems as well as achievements of the theory). There is an important take away from the theory that is *fields are inter-dependent and evolve with respect to one another*.

•Approach of this paper:

The history of Lorentz covariant theories has already taught us a lesson that *we need a concept of a 'field' in order to unify Special Relativity with Quantum Mechanics*. General Relativity is a generalization of Special Relativity. Hence, quantization of General Relativity requires a concept of a 'field' as well. It is already noted above that 'fields evolve with respect to one another'. Time is not a direct observable (in a sense that we measure time in terms of relative displacements between matter fields) in contrast to space and matter. Thus, the theory developed here is a dynamical theory of the space-matter with implicit time dependence.

ADM form of gravity is taken as a starting point which not only brings gravity closer to Lorentzian field theories but also singles out time thereby allowing to build a dynamical theory of gravity-scalar field. After separating spatial and temporal parts of fields, the full Hamiltonian $H_{\text{total}} = 0$ is reinterpreted as an equation of motion for the new combined variable field. This new field theory is then quantized canonically. Quantization of gravity and scalar field combined is a result of the quantization procedure.

2 Hamiltonian form of gravity and scalar field theory

Action for gravity together with the scalar field is taken and carried out 3+1 decomposition by foliating 4-dimensional space-time manifold into one parameter family of hypersurfaces Σ_t. Any time-like vector field T^μ decomposed into n^μ unit vector normal to hypersurface and N^μ shift vector tangential

to hypersurface. N is lapse function. (Note: $\mu, \nu..$ represents space-time indices and $a, b, c..$ represents spatial indices.)

$$T^{\mu}(t,\vec{x}) \coloneqq N(t,\vec{x})n^{\mu}(t,\vec{x}) + N^{\mu}(t,\vec{x}) \tag{1}$$

Detail discussion of 3+1 decomposition or ADM formulation can be found in references [5], section I.2.1 as well as in the book [6], section I, chapter 1.

$$\mathscr{A} = \int_{\mathbb{R}} dt \int_{M} d^3x \left(|N| \sqrt{q} \left\{ \frac{1}{16\pi G} \left(R^{(3)} + K^{ab}K_{ab} - K^2 \right) + \left(\frac{1}{2}\dot{\phi}^2 - \frac{1}{2}|\nabla\phi|^2 - V(\phi) \right) \right\} \right)(t,\vec{x}) \tag{2}$$

Curvature in 4-dimensional space $R^{(4)}$ decomposed into $R^{(3)} + K^{ab}K_{ab} - K^2$. $K_{ab}(t,\vec{x})$ is extrinsic curvature of hypersurface Σ_t, $R^{(3)}$ is a spatial curvature and $q_{ab}(t,\vec{x})$ is 3-metric both defined on 3-manifold and $\sqrt{q} = \sqrt{\det(q_{ab})}(t,\vec{x})$. Canonical conjugate momenta corresponding to $q_{ab}(t,\vec{x})$, $N^a(t,\vec{x})$, $N(t,\vec{x})$ and $\phi(t,\vec{x})$ are respectively (derived in [6], section 1.2, equation 1.2.1),

$$P^{ab}(t,\vec{x}) \coloneqq \frac{1}{16\pi G} \left(\frac{|N|}{N} \sqrt{q} \left(K^{ab} - Kq^{ab} \right) \right)(t,\vec{x}) \tag{3}$$

$$C_a(t,\vec{x}) \coloneqq \Pi_a(t,\vec{x}) = 0 \tag{4}$$

$$C(t,\vec{x}) \coloneqq \Pi(t,\vec{x}) = 0 \tag{5}$$

$$P_{\phi}(t,\vec{x}) \coloneqq \left(|N| \sqrt{q}\dot{\phi} \right)(t,\vec{x}) \tag{6}$$

$\left(q_{ab}, P^{cd}, \phi, P_{(\phi)}, N, \Pi, N^a, \Pi_a \right)(t,\vec{x})$ forms a phase space and corresponding only non-trivial Poisson Brackets (borrowed from [5], equation 1.2.1.9) are given as,

$$\left\{ q_{ab}(t,x), P^{cd}(t,x') \right\} = 16\pi G \delta^c_{(a}\delta^d_{b)}\delta^{(3)}(x,x') \tag{7}$$

$$\left\{ \phi(t,x), P_{(\phi)}(t,x') \right\} = \delta^{(3)}(x,x') \tag{8}$$

Lagrangian for gravity (refer I.2.1.7 of [5] or 1.2.5 of [6]) and for matter field is given as,

$$L_{\text{grav}} = \int_M d^3x \left\{ P^{ab}\dot{q}_{ab} + \Pi^a\dot{N}_a + \Pi\dot{N} - (\lambda C + \lambda^a C_a + N^a H_a + |N| H_{\text{scalar}}) \right\}(t,\vec{x}) \tag{9}$$

$$L_{\phi} = \int_M d^3x \left\{ P_{\phi}\dot{\phi} - |N| \sqrt{q} \left(\frac{P_{\phi}^2}{2N^2 \det(q)} + \frac{1}{2}|\nabla\phi|^2 + V(\phi) \right) \right\}(t,\vec{x}) \tag{10}$$

λ and λ^a are Lagrange undetermined multipliers and there are no equations to solve for them. Therefore, reduced constrained action (refer I.2.1.13 of [5] or 1.3.1 of [6]) is written as

$$\mathscr{A} = \int_{\mathbb{R}} dt \int_M d^3x \left\{ P^{ab}\dot{q}_{ab} + P_{\phi}\dot{\phi} - \left(N^a \mathscr{H}_a + |N| \mathscr{H}_{\text{scalar}} + |N| \mathscr{H}_{\phi} \right) \right\}(t,\vec{x}) \tag{11}$$

$$\mathscr{H}_a \coloneqq \left(-2q_{ac}D_b P^{bc} \right)(t,\vec{x}) \tag{12}$$

D_b is unique torsion-free covariant differential compatible with q_{ab}.

$$\mathscr{H}_{\text{scalar}} \coloneqq \left(\frac{16\pi G}{\sqrt{q}} \left(q_{ac}q_{bd} - \frac{1}{2}q_{ab}q_{cd} \right) P^{ab}P^{cd} - \frac{\sqrt{q}}{16\pi G} R^{(3)} \right)(t,\vec{x}) \tag{13}$$

$$\mathscr{H}_{\phi} = \sqrt{q} \left(\frac{P_{\phi}^2}{2N^2 \det(q)} + \frac{1}{2}|\nabla\phi|^2 + V(\phi) \right)(t,\vec{x}) \tag{14}$$

$\mathscr{H}_a$ are called 'Diffeomorphism constraints' or vector constraints. These constraints generate diffeomorphism on 3-Manifold. $\mathscr{H}_{\text{scalar}}$ 'Hamiltonian constraint' or scalar constraint generates temporal gauge transformation (refer 1.2.6 of [6] or I.2.1.10 of [5]). Symmetrise $\mathscr{H}_{\text{scalar}}$ in $a \leftrightarrow b$ and $c \leftrightarrow d$

$$\mathscr{H}_{\text{scalar}} = \left(-\frac{1}{2} f_{abcd} P^{ab} P^{cd} - \frac{\sqrt{q}}{16\pi G} R^{(3)}\right)(t, \vec{x}) \tag{15}$$

Where f_{abcd} is defined as,

$$f_{abcd} \coloneqq \left(\frac{16\pi G}{\sqrt{q}} \left(-q_{bc} q_{ad} - q_{ac} q_{bd} + q_{ab} q_{cd}\right)\right)(t, \vec{x}) \tag{16}$$

$$H_{\text{scalar}} = \int_M d^3x |N| \left(-\frac{1}{2} f_{abcd} P^{ab} P^{cd} - \frac{\sqrt{q}}{16\pi G} R^{(3)}\right)(t, \vec{x}) \tag{17}$$

$$H_{\text{vector}} = \int_M d^3x N^a \left(-2 q_{ac} D_b P^{bc}\right)(t, \vec{x}) \tag{18}$$

Both these constraints are first class constraints. These constraints are preserved under evolution. (Section I.2.1 of [5] can be referred for further detailed analysis) Separate time and space dependent parts of $q_{ab}(t, \vec{x})$ and $P^{ab}(t, \vec{x})$ as

$$q_{ab}(t, \vec{x}) \to q_a(t) \mathring{q}_{ab}(\vec{x}) \qquad\qquad P^{ab}(t, \vec{x}) \to P^a(t) \mathring{P}^{ab}(\vec{x}) \tag{19}$$

First term in the scalar Hamiltonain constraint becomes,

$$\begin{aligned} &\int_M d^3x |N(t)| f_{abcd}(t, \vec{x}) P^{ab}(t, \vec{x}) P^{cd}(t, \vec{x}) \\ &= |N(t)| f_{ac}(t) P^a(t) P^c(t) \int_M d^3x \left(\mathring{f}_{abcd}(\vec{x}) \mathring{P}^{ab}(\vec{x}) \mathring{P}^{cd}(\vec{x})\right) \\ &= f_{ac}(t) P^a(t) P^c(t) \end{aligned} \tag{20}$$

With,

$$\mathring{f}_{abcd}(\vec{x}) = \frac{1}{\sqrt{\mathring{q}}(\vec{x})} \left(\mathring{q}_{ab}(\vec{x}) \mathring{q}_{cd}(\vec{x}) - \mathring{q}_{ac}(\vec{x}) \mathring{q}_{bd}(\vec{x}) - \mathring{q}_{bc}(\vec{x}) \mathring{q}_{ad}(\vec{x})\right)$$

$$f_{ac}(t) \coloneqq (16\pi G)\, q_a(t) q_c(t) \tag{21}$$

$$\int_M d^3x \left(\mathring{f}_{abcd}(\vec{x}) \mathring{P}^{ab}(\vec{x}) \mathring{P}^{cd}(\vec{x})\right) = v \tag{22}$$

$$|N(t)| = \frac{\sqrt{q}(t)}{v} \tag{23}$$

Reason to choose such $|N(t)|$ is to avoid the term $\sqrt{q}(t)$ appearing in the denominator of f_{abcd}. This choice will give the kind of full Hamiltonian constraint which will allow us to construct a suitable Lagrangian for the new field which is done in the next section. Diffeomorphism constraints are given as

$$-2 \int_M d^3x N^a \left(q_{ac} D_b P^{bc}\right)(t, \vec{x}) = -2 q_i(t) P^i(t) \;\; N^a \int_M d^3x \left(\mathring{q}_{ac}(\vec{x}) D_b(\vec{x}) \mathring{P}^{bc}(\vec{x})\right) \tag{24}$$

N^a is chosen in such a way that

$$N^a \int_M d^3x \left(\mathring{q}_{ac}(\vec{x}) D_b \mathring{P}^{bc}(\vec{x})\right) = -i\hbar\,(16\pi G) \tag{25}$$

This choice of shift vector will enable us to find out Lagrangian of combined variable theory.

$$H_{\text{vector}} = 2i\hbar\,(16\pi G)\, q_i P^i \tag{26}$$

The second term in the H_{scalar} becomes

$$\frac{q(t)}{16\pi G}\frac{1}{v}\int_M d^3x\,\sqrt{\mathring{q}} R^{(3)}(t,\vec{x}) = V(\vec{q}(t)) \tag{27}$$

Although v is a function of spatial part of extrinsic curvature (because P_{ab} depends on K_{ab}), above term remains finite for $K_{ab} = 0$. Because, the first term in (17) is zero and we can choose $|N(t)| = \sqrt{q}$. That is equivalent of putting $v = 1$. By collecting (20), (26) and (27) we get gravitational part of the full Hamiltonian ('t' dependance is supressed)

$$H_{\text{scalar}} + H_{\text{vector}} = -\frac{1}{2} f_{ij} P^i P^j + 2i\hbar\,(16\pi G)\, q_i P^i - V(\vec{q}) \tag{28}$$

f_{ij} is real and symmetric matrix

$$f_{ij} = (16\pi G)\begin{pmatrix} q_1^2 & q_1 q_2 & q_1 q_3 \\ q_2 q_1 & q_2^2 & q_2 q_3 \\ q_3 q_1 & q_3 q_2 & q_3^2 \end{pmatrix} \tag{29}$$

Let us now carry out similar analysis to the matter (scalar field) part of the full Hamiltonian

$$H_{\text{matter}} = \int_M d^3x |N| \left(\frac{P_\phi^2}{2N^2\sqrt{q}} + \frac{\sqrt{q}}{2}|\nabla\phi|^2 + \sqrt{q}V(\phi)\right)(t,\vec{x}) \tag{30}$$

separate the temporal and spatial parts of the matter field and its canonical conjugate momentum $\phi(t,\vec{x}) = \phi(t)\mathring{\phi}(\vec{x})$, $P_\phi(t,\vec{x}) = P_\phi(t)\mathring{P}_\phi(\vec{x})$ and then carry out integration over spatial part we get,

$$\frac{1}{2}\int_M d^3x \frac{|N|}{N^2\sqrt{q(t)}\sqrt{\mathring{q}(\vec{x})}} P_\phi^2(t)\mathring{P}_\phi^2(\vec{x}) = \frac{vv'}{2q(t)} P_\phi^2(t) \tag{31}$$

With $v = \int_M d^3x \left(\mathring{f}_{abcd}(\vec{x})\mathring{P}^{ab}(\vec{x})\mathring{P}^{cd}(\vec{x})\right)$ and $v' = \int_M d^3x \frac{\mathring{P}^2(\vec{x})}{\sqrt{\mathring{q}}}$. Redefining $P_\phi(t)$ and $\phi(t)$ such that Poisson Bracket remain unchanged.

$$P_\phi(t) \to \sqrt{\frac{q(t)}{vv'}} P_\phi(t) \qquad\qquad \phi(t) \to \sqrt{\frac{vv'}{q(t)}}\phi(t) \tag{32}$$

$$\left\{\phi(t), P_\phi(t)\right\} = 1 \tag{33}$$

This redefinition makes

$$\text{first term of } H_{\text{matter}} = \frac{1}{2} P_\phi^2(t) \tag{34}$$

The term containing mass becomes $\mu^2\phi(t)^2$. Here, $\sqrt{q}(t)$ of lapse function cancels due to a redefinition of $\phi(t)$. $|\nabla\phi|^2$ term becomes $A\phi(t)^2$. This constant A can again be absorbed into the mass. In this manner, any coupling term can be made co-ordinate invariant by redefining coupling constants. For example ϕ^4 theory gives

$$V(\phi(t), q(t)) = \mu^2\phi(t)^2 + \lambda\frac{1}{q(t)}\phi(t)^4 \tag{35}$$

Equations (34) and (35) give matter part of the Hamiltonian

$$H_{\text{matter}} = \frac{1}{2}P_\phi^2 + V(\phi, q) \tag{36}$$

The full Hamiltonian is a sum of (28) and (36),

$$H_{\text{total}} = \frac{1}{2}P_\phi^2 - \frac{1}{2}f_{ij}P^iP^j + 2i\hbar\,(16\pi G)\,q_iP^i - V(\vec{q}) + V(\phi, q) \tag{37}$$

(q_i, P^j, ϕ, P_ϕ) is reduced phase space for H_{total} with only non-trivial Poisson brackets $\left\{q_i, P^j\right\} = \delta_i^j$ and $\left\{\phi, P_\phi\right\} = 1$. Equations of motion are given by

$$\dot{P_\phi}(t) = \left\{P_\phi(t), H_{\text{total}}\right\} \qquad \dot{\vec{P}}(t) = \left\{\vec{P}(t), H_{\text{total}}\right\} \tag{38}$$

$$\dot{\vec{q}}(t) = \{\vec{q}(t), H_{\text{total}}\} \qquad \dot{\phi}(t) = \left\{\phi(t), H_{\text{total}}\right\} \tag{39}$$

We have 3 configuration variables for gravity with one first class constraint. First class constraint reduces configuration degree of freedom by 1. These are degrees of freedom of gravity.

case 1: In special relativistic limit, $R^{(3)} = 0$ imply $V(\vec{q}) =$ const. and extrinsic curvature ($K_{ab} = 0$) imply $P^j = 0$. Then,

$$H_{\text{total}} = \frac{1}{2}P_\phi^2 + V(\phi)$$

Note that $V(\phi, q) = V(\phi)$ because $q(t) = 1$. This is exactly the Hamiltonian for scalar field with a spatial part being integrated over and contribution of that part is absorbed inside coupling constants. This Hamiltonian gives time evolution.

case 2: In zero scalar field limit, $V(\phi, q) = 0$. Equation of motion give $\dot{\phi} = P_\phi$ but since $\phi = 0$ and $\dot{\phi} = 0 \rightarrow P_\phi = 0$. Therefore,

$$H_{\text{total}} = -\frac{1}{2}f_{ij}P^iP^j + 2i\hbar\,(16\pi G)\,q_iP^i - V(\vec{q})$$

This is the Hamiltonian for gravity with a lapse function and a shift vector chosen appropriately. Hamiltonian does not give physical time evolution. Instead, it gives a gauge transformation.

3 Combined variable theory of gravity and scalar field

3.1 Classical Theory

Rewrite (37) by using $P_\phi = -i\hbar\frac{\partial}{\partial\phi} = -i\hbar\partial_\phi$, $P^k = -i\hbar\frac{\partial}{\partial q_k} = -i\hbar\partial^k$. $\frac{\hbar^2}{2}V(q_k,\phi) \coloneqq V(\vec{q}) - V(\phi,q)$ is chosen only for convenience.

$$\hat{H}_{\text{total}}\Phi = \left(-\frac{\hbar^2}{2}\frac{\partial^2}{\partial\phi^2} + \frac{\hbar^2}{2}f_{ij}\frac{\partial}{\partial q_i}\frac{\partial}{\partial q_j} + 2\hbar^2\,(16\pi G)\,q_i\frac{\partial}{\partial q_j} - \frac{\hbar^2}{2}V(q_k,\phi)\right)\Phi$$

[Note:'t' dependence is suppressed from here onwords.] This equation is now interpreted $\hat{H}_{\text{total}}\Phi = 0$ as an equation of motion for classical field Φ of combined variables.

$$-\frac{\hbar^2}{2}\left(\frac{\partial^2}{\partial\phi^2} - \frac{\partial}{\partial q_i}f_{ij}\frac{\partial}{\partial q_j} + V(q_k,\phi)\right)\Phi = 0 \tag{40}$$

$\frac{\partial}{\partial q_i}f_{ij}\frac{\partial}{\partial q_j} = f_{ij}\frac{\partial}{\partial q_i}\frac{\partial}{\partial q_j} + \frac{\partial f_{ij}}{\partial q_i}\frac{\partial}{\partial q_j}$. In the second term, $\frac{\partial}{\partial q_i}$ acting on q_i of f_{ij} gives $3q_j$ and acting on q_j of f_{ij} gives $1\,q_j$. This second term is a result of Diffeomorphism constraints. If we had chosen a different shift vector, that would have introduced extra derivative coupling but the choice (25) compensates this extra derivative term. Taking plane wave solution $\Phi \sim e^{i(P_\phi\phi + P^k q_k)}$ we recover (37) which has special relativistic as well as zero scalar field limit.

$\vec{q}$ is dimensionless and ϕ is made dimensionless by redefining $\phi \to \frac{1}{\sqrt{16\pi G}}\phi$. Also by absorbing $(16\pi G)$ inside $V(q_k,\phi)$ we get

$$\left(\partial_\phi^2 - \partial^i\eta_{ij}\partial^j + V(q_k,\phi)\right)\Phi(\phi,q_k) = 0 \tag{41}$$

$\eta_{ij} \coloneqq q_i q_j$ will serve as a spatial metric for the space of $(\phi,\vec{q})$. Action for this combined variable field is given as

$$\mathscr{A} = \int d\phi \int d^D q\left\{\frac{1}{2}(\partial_\phi\Phi)^2 - \frac{1}{2}(\partial^i\Phi)\eta_{ij}(\partial^j\Phi) - \frac{1}{2}V(q_k,\phi)\Phi^2\right\} \tag{42}$$

$D = 0, 1, 2$ or 3 depending on how many components of $\vec{q}$ are time dependent. $V(q_k,\phi)$ can be positive or negative. For $V < 0$ and $\alpha > 0$

$$\mathscr{A} = \int d\phi \int d^D q\left\{\frac{1}{2}(\partial_\phi\Phi)^2 - \frac{1}{2}(\partial^i\Phi)\eta_{ij}(\partial^j\Phi) - \frac{1}{2}V\Phi^2 - \frac{1}{4}\alpha\Phi^4\right\} \tag{43}$$

The field is unstable at $\Phi = 0$ and will condense into a stable state $\Phi\,|_{\text{vac}} = \pm\sqrt{\frac{-V}{\alpha}}$. Symmetry of the ground state is spontaneously broken. If we shift $\Phi = \rho - \Phi\,|_{\text{vac}}$, we get

$$-\frac{1}{2}V\Phi^2 - \frac{1}{4}\alpha\Phi^4 \longrightarrow \frac{1}{2}2V\rho^2 - \frac{1}{4}\alpha\rho^4 + \left(V\Phi\,|_{\text{vac}} + \alpha\Phi\,|_{\text{vac}}^3\right)\rho + \alpha\Phi\,|_{\text{vac}}\,\rho^3 \tag{44}$$

Extra constant terms are not written as they are inconsequential. ρ^2 term comes with a correct sign and therefore mass spectrum remain positive definite. Treatment of this theory is beyond scope of this paper and will be discussed in

the next paper.

A 4-dimensional space-matter manifold is equipped with a metric

$$g_{\mu\nu} \coloneqq \begin{pmatrix} 1 & 0 \\ 0 & -\eta_{ij} \end{pmatrix} \tag{45}$$

which allows us to measure the length in the space of $(\phi, \vec{q})$

$$dl^2 = g_{\mu\nu} dq^\mu dq^\nu = d\phi^2 - \eta_{ij} dq^i dq^j \tag{46}$$

Rewrite action (42) in covariant form

$$\mathscr{A} = \int d\phi \int d^D q \left\{ \frac{1}{2} g_{\mu\nu} \partial^\mu \Phi \partial^\nu \Phi - \frac{1}{2} V(q_k, \phi) \Phi^2 \right\} \tag{47}$$

$\mu, \nu = 0, 1, 2, 3$ with 0^{th} term being ϕ index and $1, 2, 3$ are metric indices. $\partial^\mu = \frac{\partial}{\partial q_\mu} = \left(\frac{\partial}{\partial \phi}, \frac{\partial}{\partial q_i} \right)$. Define

$$\Pi_\mu \coloneqq \frac{\partial \mathscr{L}}{\partial(\partial^\mu \Phi)} = g_{\mu\nu} \partial^\nu \Phi \tag{48}$$

Extremizing action under infinitesimal variation of combined variable field $\Phi \to \Phi + \delta\Phi$ we get

$$- \int d\phi \int d^D q \left(\partial^\mu g_{\mu\nu} \partial^\nu \Phi + V(q_k, \phi) \Phi \right) \delta\Phi + \int d\phi \int d^D q \partial^\mu \left(g_{\mu\nu} \partial^\nu \Phi \, \delta\Phi \right) = 0 \tag{49}$$

Second term is a surface term and if variations are such that $\delta\Phi \to 0$ on the surface then we get (41). Equivalently, current ($J_\mu \coloneqq g_{\mu\nu} \partial^\nu \Phi \, \delta\Phi$) is conserved if equation of motion is satisfied. Invariance of action under $q_\mu \to q_\mu + \delta q_\mu$ results into

$$\partial^\rho T^\mu_\rho = \partial^\rho \left(\frac{\partial \mathscr{L}}{\partial(\partial^\rho \Phi)} \partial^\mu \Phi - \mathscr{L} \delta^\mu_\rho \right) = 0$$

This gives us four constants of motion. $\mu = 0$ gives energy and other three are momentum of the combined variable field

$$E = \int d^D q \left(\frac{\partial \Phi}{\partial \phi} \frac{\partial \Phi}{\partial \phi} + \frac{\partial \Phi}{\partial q_i} \eta_{ij} \frac{\partial \Phi}{\partial q_j} + V(q_k, \phi) \Phi^2 \right) \tag{50}$$

$$\mathbf{P}_i = \int d^D q \;\; \frac{\partial \Phi}{\partial \phi} \frac{\partial \Phi}{\partial q_i} \tag{51}$$

Hamiltonian is obtained by carrying out Legendre transformation

$$\mathbf{H} = \int d^D q \left(\frac{1}{2} \Pi^2 + \frac{1}{2} \frac{\partial \Phi}{\partial q_i} \eta_{ij} \frac{\partial \Phi}{\partial q_j} + \frac{1}{2} V(q_k, \phi) \Phi^2 \right) \tag{52}$$

Π ($\Pi_0 = \frac{\partial \Phi}{\partial \phi}$ of (48)) is a canonical conjugate momentum corresponds to the field Φ.

$$\begin{aligned} &\left\{ \Phi(\phi, \vec{q}), \Pi(\phi, \vec{q}') \right\} \coloneqq \delta(\vec{q}, \vec{q}') \\ &\left\{ \Phi(\phi, \vec{q}), \Phi(\phi, \vec{q}') \right\} \coloneqq 0 \qquad \left\{ \Pi(\phi, \vec{q}), \Pi(\phi, \vec{q}') \right\} \coloneqq 0 \end{aligned} \tag{53}$$

Dynamics of the theory is given by following equations of motion.

$$\frac{\partial \Phi}{\partial \phi} = \{\Phi, \mathbf{H}\}\,, \qquad \frac{\partial \Pi}{\partial \phi} = \{\Pi, \mathbf{H}\} \tag{54}$$

Since,

$$\frac{\partial \Phi(\vec{q}')}{\partial \phi} = \{\Phi(\vec{q}'), \mathbf{H}\} = \int d^D q\, \Pi(\vec{q}) \delta(\vec{q}, \vec{q}') = \Pi(\vec{q}')$$

evolution is consistent. The second equation gives evolution of Π with ϕ.

•**Remarks:**

In order to see physics of classical combined variable field theory, let us first revisit a transition

$$\text{particle} \rightarrow \text{field}$$

I. $H = \frac{p^2}{2m} + V(\vec{x}, t)$ is the Hamiltonian for a particle or system of particles. Hamiltonian equations give trajectory (path) of a particle or system of particles.

II. $\hat{H}\phi = \left(\frac{\hat{p}^2}{2m} + \hat{V}(\vec{x}, t)\right)\phi$ is reinterpreted as an equation of motion for classical field which is defined over space and evolves with time. The second term is seen as coupling which is $(\vec{x}, t)$ dependent.

Now, examine a transition

$$\text{field} \rightarrow \text{combined variable field}$$

III. In the case of gravity, space-time itself is a dynamical entity. Hamiltonian equations of motion tell us how ϕ varies with 3-metric or equivalently how q_i changes with ϕ.

IV. Unlike case III, combined variable field Φ is spread over the space of $\vec{q}(t)$. Hamiltonian (52) gives ϕ evolution. 3-curvature, spatial part of extrinsic curvature (through v) and scalar field couplings contribute to the potential energy of the combined variable field.

3.2 Quantum Theory

Theory is quantized by raising Poisson brackets of combined variables to commutators

$$\left[\hat{\Phi}(\phi, \vec{q}), \hat{\Phi}(\phi, \vec{q}')\right] = \left[\hat{\Pi}(\phi, \vec{q}), \hat{\Pi}(\phi, \vec{q}')\right] = 0 \tag{55}$$
$$\left[\hat{\Phi}(\phi, \vec{q}), \hat{\Pi}(\phi, \vec{q}')\right] = i\sqrt{\frac{h}{G}}\delta(\vec{q}, \vec{q}')$$

(50) and (48) shows that Φ and Π both have dimensions of $\sqrt{E}$. Therefore, dimensional constant (Planck energy) $\sqrt{\frac{h}{G}}$ is introduced. Also, define creation and annihilation operators

$$\hat{a} := \frac{1}{\sqrt{2}}\left(\hat{\Pi} - i\omega\hat{\Phi} - i\mathcal{L}_{\vec{q}}\hat{\Phi}\right), \qquad \hat{a}^\dagger := \frac{1}{\sqrt{2}}\left(\hat{\Pi} + i\omega\hat{\Phi} + i\mathcal{L}_{\vec{q}}\hat{\Phi}\right) \tag{56}$$

$\omega(\phi, \vec{q})$ is a solution of (60). $\mathcal{L}_{\vec{q}} := \vec{q}.\frac{\vec{\partial}}{\partial q}$ is directional derivative. $\mathcal{L}_{\vec{q}}\hat{\Phi}$ gives variation of Φ along $\vec{q}$.

$$\begin{aligned}\hat{a}^\dagger\hat{a} = \frac{1}{2}\hat{\Pi}^2 + \frac{1}{2}\left(\mathcal{L}_{\vec{q}}\hat{\Phi}\right)^2 + \frac{1}{2}\omega^2\hat{\Phi}^2 + i\frac{1}{2}\omega\left[\hat{\Phi}, \hat{\Pi}\right] \\ -i\frac{1}{2}\left[\hat{\Pi}, \mathcal{L}_{\vec{q}}\hat{\Phi}\right] + \omega\left(\mathcal{L}_{\vec{q}}\hat{\Phi}\right)\hat{\Phi}\end{aligned} \tag{57}$$

Notice that the second term in above equation is exactly the second term in the Hamiltonian (52). Since,

$$\left[\hat{\Pi}(\vec{q}), \mathcal{L}_{\vec{q}'}\hat{\Phi}(\vec{q}')\right] = \sqrt{\frac{h}{G}}\mathcal{L}_{\vec{q}'}\delta(\vec{q}, \vec{q}') \tag{58}$$

The forth term in (57) becomes $\mathcal{L}_{\vec{q}}\delta(\vec{0}) = 0$. Let us now calculate the last term

$$\int d^D q\ \omega\widehat{\left(q_i\frac{\partial\Phi}{\partial q_i}\right)}\hat{\Phi} = \int d^D q \frac{\partial}{\partial q_i}.\left(\frac{1}{2}\omega q_i\hat{\Phi}^2\right) - \int d^D q\left(\frac{\vec{\partial}}{\partial q}.\left(\frac{1}{2}\vec{q}\omega\right)\right)\hat{\Phi}^2 \tag{59}$$

If variable Φ is chosen such that $\lim_{q\to\pm\infty}\Phi \to 0$ then total derivative term vanishes and (57) becomes

$$\hat{a}^\dagger\hat{a} = \frac{1}{2}\hat{\Pi}^2 + \frac{1}{2}\left(\mathcal{L}_{\vec{q}}\hat{\Phi}\right)^2 + \frac{1}{2}\left(\omega^2 - \frac{\vec{\partial}}{\partial q}.(\vec{q}\omega)\right)\hat{\Phi}^2 - \frac{1}{2}\sqrt{\frac{h}{G}}\omega\delta(\vec{0})$$

In order to get the Hamiltonian we set

$$V = \omega^2 - \frac{\vec{\partial}}{\partial q}.(\vec{q}\omega) = \omega^2 - D\omega - \vec{q}.\frac{\partial\omega}{\partial\vec{q}} \tag{60}$$

$\omega(\phi, \vec{q})$ is a solution to the first order differential equation.

$$\hat{a}^\dagger\hat{a} = \frac{1}{2}\hat{\Pi}^2 + \frac{1}{2}\left(\mathcal{L}_{\vec{q}}\hat{\Phi}\right)^2 + \frac{1}{2}V\hat{\Phi}^2 - \frac{1}{2}\sqrt{\frac{h}{G}}\ \omega(\phi, \vec{q})\ \delta(\vec{0})$$

Then Hamiltonian operator can be written as

$$\hat{\mathbf{H}} = \int d^D q\left(\hat{a}^\dagger\hat{a} + \frac{1}{2}\sqrt{\frac{h}{G}}\ \omega(\phi, \vec{q})\ \delta(\vec{0})\right) \tag{61}$$

Commutation relations of creation and annihilation operators can be obtained by using (55), (56) and (58). Only non-trivial commutation relation

$$\left[\hat{a}(\vec{q}), \hat{a}^\dagger(\vec{q}')\right] = \sqrt{\frac{h}{G}}\left(\omega(\phi, \vec{q}) + \mathcal{L}_{\vec{q}}\right)\delta(\vec{q}, \vec{q}')$$

But $\mathcal{L}_{\vec{q}}\delta(\vec{q},\vec{q}') = \vec{q}.\frac{\vec{\partial}}{\partial q}\delta(\vec{q},\vec{q}') = \sum_i q_i \frac{\partial}{\partial q_i}\delta(\vec{q},\vec{q}')$ and since $x\frac{\partial}{\partial x}\delta(x) = -\delta(x)$ implying $\mathcal{L}_{\vec{q}}\delta(\vec{q},\vec{q}') = -D\delta(\vec{q},\vec{q}')$.

$$\left[\hat{a}(\vec{q}),\hat{a}^\dagger(\vec{q}')\right] = \sqrt{\frac{h}{G}}\left(\omega(\phi,\vec{q}) - D\right)\delta(\vec{q},\vec{q}') \tag{62}$$

Rewriting Hamiltonian in terms of number operator $\hat{n}$ which gives number of field quantum.

$$\hat{\mathbf{H}} = \int d^D q\ \sqrt{\frac{h}{G}}\left(\omega(\phi,\vec{q}) - D\right)\hat{n} + \int d^D q \frac{1}{2}\sqrt{\frac{h}{G}}\ \omega(\phi,\vec{q})\ \delta(\vec{0}) \tag{63}$$

•**Remarks:**
The state $|n,\vec{q}>$ is an eigen state of the Hamiltonian operator. $\hat{\Phi}(\vec{q})$ is a sum of creation and annihilation operators, acting on the vacuum produces a field quanta with 3-metric $\vec{q}$. A field quanta (or a particle) is seen as an excitation in the combined variable field. The second term is vacuum energy term. A role of creation and annihilation operator get reversed when $(\omega - D) < 0$. For example, in the case of $\omega(\phi,\vec{q}) = 0$

$$\left[\hat{a}^\dagger(\vec{q}),\hat{a}(\vec{q}')\right] = \left(D\sqrt{\frac{h}{G}}\right)\delta(\vec{q},\vec{q}') \tag{64}$$

$$\hat{\mathbf{H}} = \int d^D q\left(D\sqrt{\frac{h}{G}}\right)\hat{n} \tag{65}$$

Homogeneous and non-stationary space ($q_{ab}(t,\vec{x}) = q_{ab}(t)$): Since, $\partial_a q_{bc} = 0$ implying spatial curvature tensor is 0 and $R^{(3)}(q_{ab},\partial_a q_{bc}) = 0 \rightarrow \omega(\vec{q},\phi) = \omega(\phi)$. (60) reduces to algebraic equation.

Inhomogeneous and stationary space ($q_{ab}(t,\vec{x}) = q_{ab}(\vec{x})$): Since 3-metric is time independent, second term in the (52) vanishes. Because action functional in this case is $q(t)$ independent. Lagrangian and Hamiltonian does not invlove integration over q.

$$\mathbf{H} = \frac{1}{2}\Pi^2 + \frac{1}{2}V(v,R^{(3)},\phi)\ \Phi^2 \tag{66}$$

(60) reduces to $\omega^2 = V$. Treatment for negative V is suggested in (44). Hamiltonian operator for this theory is

$$\hat{\mathbf{H}} = \sqrt{\frac{h}{G}}\ \omega(\phi,\vec{q})\ \hat{n} + \frac{1}{2}\sqrt{\frac{h}{G}}\ \omega(\phi,\vec{q})\ \delta(\vec{0}) \tag{67}$$

4 Acknowledgment

I am immensely grateful to Prof. Ajay Patwardhan for providing expertise. This work would not have possible without his guidance. Any errors are my own and should not tarnish his reputation.

5 Bibliography

[1] The Dynamics of General Relativity - R. Arnowitt, S. Deser, C. W. Misner. arXiv:gr-qc/0405109v1 19 May 2004.

[2] Background Independent Quantum Gravity: A status report - A. Ashtekar, J. Lewandowsky. arXiv:gr-qc/0404018v2 3 Sept 2002.

[3] Gravity and the Quantum - A. Ashtekar. arXiv:gr-qc/0410054v2 19 Oct 2004

[4] Loop quantum gravity: the first twenty five years - Carlo Rovelli. arXiv:1012.4707v5 [gr-qc] 28 Jan 2012

[5] Lectures on Loop Quantum Gravity - T. Thiemann arXiv:gr-qc/0210094v1 28 Oct 2002

[6] Modern Canonical Quantum General Relativity - T. Thiemann Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, U.K., 2007.